SERVICES OFFERED

Family Planning-

Health services to assist women in preventing or achieving pregnancy. All medically accepted methods of birth control available.

Clinical Services-

Available to men and women, includes physical exam, STI/HIV testing, cervical cancer screening and testing for UTI.

Wisconsin Well Woman Program

Emergency Contraception

Pregnancy Testing

Prenatal Care Coordination

A sliding-fee payment scale is available and is based on family size and income.

Please call to schedule an appointment.



PIERCE COUNTY REPRODUCTIVE HEALTH

1234 S Wasson Lane, Suite A PO Box 82 River Falls, WI 54022

Phone: 715-425-8003

Website: http://www.co.pierce.wi.us/Public%20Health/PH-New-Pregnancy.html "Like" us on Facebook @
Pierce County Public Health Department

PIERCE COUNTY REPRODUCTIVE HEALTH

Men's health online resources

Telephone: 715-425-8003

BIRTH CONTROL

For information available on all types of birth control methods:

www.wisconsinfamilyplanning.org

www.womenshealth.gov

www.webmd.com

www.arhp.org/MethodMatch

www.contraceptivetechnology.org

For information regarding specific types of birth control:

Mirena IUD

www.mirena-us.com

Implanon

www.implanon-usa.com

ParaGard IUD www.paragard.com

Nuva Ring www.nuvaring.com

Plan B/Emergency Contraception

www.not-2-late.com

www.planbonestep.com

Men's Health Online Resources

SEXUALLY TRANSMITTED INFECTIONS

Nearly half of all new STD cases are in young people between the ages of 15 and 24.*

In 2009, young persons (ages 13-29) accounted for 39% of all new HIV infections in the US.*

There were 1.4 million new cases of chlamydia in 2011, people aged 15-24 had 4 times the reported rate of the total population.*

http://www.itsyoursexlife.com/gyt/

http://www.cdc.gov/std/default.htm

http://www.nlm.nih.gov/medlineplus/sexuallytransmitteddiseases.

http://www.nlm.nih.gov/medlineplus/tutorials/sexuallytransmitteddiseases/htm/_no_50_no_0.htm

MENTAL HEALTH

Every 13.7 minutes someone in the United States dies by suicide.**

90% of people who die by suicide have a diagnosable and treatable psychiatric disorder at the time of their death.**

Males between ages 15-19 are five times more likely, and between ages 20-24 seven times more likely than females to commit suicide. Over 80% of suicide deaths are men.***

http://www.suicidepreventionlifeline.org/

http://www.webmd.com/mental-health/recognizing-suicidal-behavior

ALCOHOL ABUSE

Men are more likely than women to drink excessively and take other risks (e.g., drive fast or without a safety belt), increasing their risk of injury or death.*



Men consistently have higher rates of alcohol-related deaths and hospitalizations than women.*

Alcohol use by men increases the chances of engaging in risky sexual activity including unprotected sex, sex with multiple partners, or sex with a partner at risk for sexually transmitted diseases.*

http://www.cdc.gov/alcohol/fact-sheets/mens-health.htm

http://www.goerie.com/substanceabuse/risks_rise_when_teens_combine_.html

GAY AND BISEXUAL MEN

The rate of new HIV diagnoses among men who have sex with men (MSM) is more than 44 times that of other men.*

The rate of primary and secondary syphilis among MSM is more than 46 times that of other men and more than 71 times that of women.*

http://blog.aids.gov/2012/12/syphilis-and-hiv-a-dangerous-duo-affecting-gay-and-bisexual-men.html

http://www.cdc.gov/hiv/topics/msm/index.htm

http://www.gmhc.org

*Centers for Disease Control and Prevention, 2012.

**American Foundation for Suicide Prevention, 2013

***Men's Health Network, (n.d.).



